




# Asbestos and other occupational lung diseases in New Zealand

2010 ANNUAL REPORT



DOL-11670 MAR 11



**Front Cover Picture:** *Anthrophyllite asbestos, viewed through a scanning electron microscope.*

From <http://usgsprobe.cr.usgs.gov/picts2.html>. Image contains materials that came originally from the United States Geological Survey, an agency of the United States Department of Interior. Sourced from Wikimedia [www.wikimedia.org](http://www.wikimedia.org).

Disclaimer: The Department of Labour has made every effort to ensure that the information contained in this publication is reliable, but makes no guarantee of its completeness. The Department may change the contents of this guideline at any time without notice.

© Crown copyright 2011

This material is Crown copyright unless otherwise stated and may be reproduced free of charge without requiring specific permission. This is subject to it being reproduced accurately and not being used in a derogatory manner or in a misleading context. The source and copyright status should be acknowledged. The permission to reproduce Crown copyright protected material does not extend to any material in this report that is identified as being the copyright of a third party.

Published by the Department of Labour  
PO Box 3705, Wellington 6140, New Zealand  
[www.dol.govt.nz](http://www.dol.govt.nz)  
March 2011

ISBN: 978-0-478-36023-3

# CONTENTS

<b>CONTENTS .....</b>	<b>3</b>
<b>INTRODUCTION .....</b>	<b>4</b>
<b>SUMMARY OF CHANGES.....</b>	<b>5</b>
<b>PART 1: ASBESTOS .....</b>	<b>7</b>
1.1 What is Asbestos? .....	7
<b>PART 2: REVIEW OF ASBESTOS-RELATED DISEASE NOTIFICATIONS.....</b>	<b>13</b>
2.1 Summary.....	13
2.2 Results .....	14
2.3 Mesothelioma .....	15
2.4 National Cancer Figures for Mesothelioma.....	17
2.5 Lung Cancer .....	18
2.6 Asbestosis .....	19
2.7 Pleural Abnormalities.....	21
2.8 Discussion .....	22
<b>PART 3: NON-ASBESTOS OCCUPATIONAL RESPIRATORY DISEASE.....</b>	<b>24</b>
3.1 Occupational Asthma .....	24
3.2 Occupational Asthma Case Studies .....	25
3.3 Metal Fume Fever .....	27
3.4 Chronic Obstructive Pulmonary Disease.....	27
3.5 Silica and Silicosis – One Agent, Many Outcomes.....	28
3.6 Indoor Air Quality (IAQ).....	28
3.7 Work-Related Occupational Respiratory Disease .....	29
3.8 Other asbestos issues.....	29
<b>APPENDIX 1 .....</b>	<b>31</b>
Contact details for the Asbestos and Occupational Respiratory Disease Registrar .....	31
Members of the National Asbestos Medical Panel.....	31
Obituary – A H Grootegoed.....	31
<b>APPENDIX 2 .....</b>	<b>32</b>
The National Asbestos Registers .....	32
<b>APPENDIX 3 .....</b>	<b>37</b>
Department of Labour Publications on Asbestos.....	37
<b>APPENDIX 4 .....</b>	<b>39</b>
References.....	39

## INTRODUCTION

The Department of Labour's two Asbestos Registers - the *Disease Register* and the *Exposure Register* - have been in existence since 1992. When established, they had the following objectives:

1. to raise awareness of asbestos-related disease nationally;
2. to improve the radiological diagnosis of asbestos disease by using the ILO reference radiographs;
3. to confirm that pleural plaques is a disease and not just a marker for asbestos exposure; and
4. to reinforce the importance of patient support groups.

The legacy of disease from working with and exposure to asbestos in New Zealand is continuing. Exposure is now largely confined to brake repairers, lino removers, demolition workers, carpenters, electrical, plumbing and building maintenance workers, and asbestos removal workers.

Worldwide, asbestos exposure still occurs at a high level; and there are a number of lessons than can be learned as a result.

Firstly, there continues to be a general lack of awareness of the effects of inhaling dust at work, regardless of the type of dust. Secondly, too great an emphasis has been put on whether or not dust is a danger to health. Concepts that cement dust is safe and silica dust is unsafe; or blue asbestos is worse than white asbestos, have distracted governments from the real issue of dirty workplace air.

Today, this is recognised as a major problem, both in terms of respiratory and general health. However, its solutions are not complex. They involve putting in place good general and local exhaust ventilation, and providing comfortable and effective respiratory protection. While that sounds relatively simple, there needs to be a desire for clean workplace air.

With a gradual reduction in cigarette smoking among the working population, the impact of dirty workplace air on respiratory health is likely to become clearer, with diseases such as welder's lung, asthma, and chronic obstructive pulmonary disease recognised more clearly for their relationship to work practices and procedures.

Dr W I Glass (Convenor)  
Dr R Armstrong  
Dr D Jones  
Dr A Brant

**National Asbestos Medical Panel**

March 2011

## SUMMARY OF CHANGES

Compared with the [previous year's](#) annual report:

Cases	1992-2008	1992-2010	Increase
Mesothelioma	212	224	12
Lung cancer	107	116	9
Asbestosis	253	269	16
Pleural Abnormalities	553	589	36
<b>Total</b>	<b>1125</b>	<b>1198</b>	<b>73</b>

**Table 1: Categories of Notified Occupational Lung Diseases**

Criteria	1992-2008	1992-2010	Increase
Caucasian	204	215	11
Maori	6	No change	-
Other	2	3	1
Males	199	210	11
Females	13	14	1
Current Smokers	19	No change	-
Ex-Smokers	116	125	9
Never Smoked	59	No change	-

**Table 2: Mesothelioma Data**

Year	Number
2003	78
2004	82
2005	102
2006	80
2007	91

**Table 3: National Cancer Figures for Mesothelioma**

Criteria	1992-2008	1992-2010	Increase
Caucasian	101	110	9
Maori	4	No change	-
Pacific Islander	1	No change	-
Other	1	No change	-
Males	105	114	9
Females	2	No change	-
Current Smokers	25	No change	-
Ex-Smokers	72	81	9
Never Smoked	7	No change	-

**Table 4: Lung Cancer Data**

Criteria	1992-2008	1992-2010	Increase
Caucasian	249	265	16
Maori	2	No change	-
Pacific Islander	2	No change	-
Males	237	253	16
Females	16	No change	-
Current Smokers	20	21	1
Ex-Smokers	183	192	9
Never Smoked	43	49	6

**Table 5: Asbestosis Data**

Criteria	1992-2008	1992-2010	Increase
Caucasian	542	577	35
Maori	7	8	1
Pacific Islander	4	No change	-
Males	543	579	36
Females	10	No change	-
Current Smokers	52	59	7
Ex-Smokers	328	349	21
Never Smoked	138	148	10


**Table 6: Pleural Abnormalities Data**

## PART 1: ASBESTOS

### 1.1 What is Asbestos?

Asbestos is a term used to describe naturally occurring fibrous hydrated silicates of which there are six common varieties.

Figure 1 below shows the different types of asbestos.



**Figure 1: Types of Asbestos**

Amosite (brown asbestos) and crocidolite (blue asbestos) have straight needle-like fibres which naturally split in their long axis producing very fine fibrils (thin fibres). In contrast, chrysotile (white asbestos) has softer curlier hair-like fibres, which makes them useful for weaving and have been used to make asbestos cloth for centuries.

When these fibres are inhaled they behave in different ways in the lungs.

Chrysotile (magnesium silicate) fibres slowly lose magnesium, which leaches into body fluids making the fibre more easily digested by scavenger cells (macrophages). This reduces the body burden of the asbestos dust, and has led to the view that white fibres are less harmful.

Crocidolite (iron-sodium silicate) fibres are straight and rigid and can split longitudinally. They are more resistant to body fluids and can survive unchanged in the body for up to 40 years or more. The fine fibrils can migrate through the lung tissue to the lung lining (pleura).

Amosite (iron magnesium silicate) fibres are longer, making it useful for insulation. Again, the fibres remain unchanged in the body for years.

### **1.1.1 Asbestos awareness in New Zealand**

**1938** The introduction to the *Report of the Interdepartmental Committee on Silicosis* refers to asbestos as follows:

*"In the working of asbestos the dust produced gives rise to a pulmonary condition known as asbestosis, which, although different in its character from silicosis, causes a disease similar in some respects to it. Free silica is not associated with asbestos, yet the fibres produce a deadly pulmonary disease."*

**1951** In the *Annual Report of the Department of Health* Dr Garland, Director of the Division of Occupational Health, Notes:

*"Asbestos is now being quarried in the Dominion, and the dust can be expected to cause a certain amount of lung damage unless proper precautions are taken."*

**1953** Notification of occupational diseases introduced. No specific identification of asbestos-related lung disease.

**1960** *Occupational Health Notes on Diseases Arising from Occupation* published by the Department of Health:

*"The fibrosis of asbestos is diffuse ...there is some statistical evidence for an increased incidence of bronchogenic carcinoma."*

**1964** A New Zealand standard is set for asbestos fibres in air.

**1968** Dr Copplestone, Assistant Director (Occupational Health) initiates a New Zealand asbestos survey.

**1971** Dr Allingham, who succeeded Dr Copplestone, reports on the survey:

*"207 workers surveyed. 101 workers x-rayed. 17 showed pleural changes (one case later confirmed as asbestosis)."*

**1977** Environmental Dust Laboratory established to identify asbestos fibres in air.

**1978** Asbestos Regulations promulgated.

**1981** The allowable concentration for asbestos fibres (other than crocidolite fibres) was reduced from 2 fibres/ml to 1 fibre/ml.

Crocidolite remained at 0.2 fibres/ml over a 10-minute sampling period.

- 1982** Jim Butterworth, Auckland Secretary, NZ Engineers Union, expresses concern about long-term effects of asbestos on workers at James Hardie factory.
- 1983** *Interim Report of Workers at James Hardie* (unpublished)  
Kjellstrom T and Rennie F. Of 353 workers x-rayed, 53 showed parenchymal and/or pleural changes. An increasing percentage of abnormalities found over time since first exposure.
- Asbestos Regulations replace 1978 regulations.
- A *New Zealand Gazette* notice reduced air standard for crocidolite to 0.1 fibre/ml over 4 hours.
- 1984** *Relative Cancer Risks from Exposure to Different Asbestos Fibre Types*. J. Keir Howard, *NZ Medical Journal* 97; 646-9. In this review article, the point is made that:
- "...considerable amounts of crocidolite were used in the past, particularly in insulation, and as New Zealand did not restrict the use of the amphiboles until well after most industrialised nations had done so, the use of blue asbestos continued for longer in this country."*
- Importation of blue and brown asbestos banned in New Zealand.
- 1987** Asbestos Regulations 1983, Amendment No. 1, tightens provisions for medical examinations.
- 1988** Thelma Bell, widow of asbestosis victim Clarrie, organises a public meeting in Christchurch.
- Barry Brown, Secretary of the South Island Labourers Union, arranges for Maevis Watson, Occupational Health Nurse and Associate Professor Bill Glass from the Otago Medical School, to interview ex-Fletcher workers from the Mandeville Street plant.
- Cate Brett – *Christchurch Star* – highlights the tragedy of asbestos use.
- 1989** Robin McKenzie, Engineer with NZED (New Zealand Electricity Department), is diagnosed as suffering from mesothelioma. In 1990, his lawyer took legal action against the Crown. Media coverage keeps up public concern.

- 1990** *Review of Fletcher's Mandeville Street Factory* (unpublished) Glass W.I. 103 workers contacted. 87 investigated. 47 cases of asbestos-related conditions (plus 7 probable cases). 12 deaths to date.
- Hon. Bill Birch, Minister of Labour, establishes the Asbestos Advisory Committee in October to report on all aspects of asbestos exposure in New Zealand.
- 1991** *Lung Cancer, Smoking and Exposure to Asbestos in New Zealand.* Glass W.I; Kawachi I., Pearce N. *Journal of Occupational Health Safety* 7 (1).
- "Case control studies based on NZ Cancer Registry showed that asbestos-related occupations were found to be associated with elevated risks of cancer in the lung, pleura and peritoneum. The risks of cancer in these three sites were highest among the group comprising machinery fitters, plumbers, welders, boilermakers, metal moulders, metal polishers and electricians."*
- Report of the Asbestos Advisory Committee* presented to the Minister in April and Minister decides to implement the recommendations following Cabinet approval in August.
- Tim Frederiksen establishes the Asbestos Victim and Support Group, which later became the Asbestos Disease Association of New Zealand.
- 1992** In March, a National Asbestos Register was established (recommendation 4 of the Report). This Register was to be in two parts: Part 1 for those who had been exposed and Part 2 for those suffering from an asbestos-related disease. The Occupational Safety and Health Service of the Department of Labour was appointed to administer the registers.
- ECNZ (Electricity Corporation of New Zealand) combines with the Department of Labour to publicise asbestos hazards.
- An audit of floor sanders and work practices involving asbestos-backed vinyl sheeting in the Christchurch area was carried out by KD Sheat and published by OSH (Department of Labour).
- ACC lump sum entitlements for asbestos-related claims ceased.
- 1993** The first *Annual Report of the National Asbestos Registers* confirms 199 cases of asbestos-related disease.

(March) Rights to initiate common law claims in regard to asbestos-related conditions ceased.

ECNZ establishes model asbestos surveillance programme of current and past employees.

Secondary cases of asbestos disease found to occur in family members of asbestos workers whose only exposure was to dust, brought to the home from the workplace.

- 1994** The second Annual Report reviews and confirms a total of 462 cases of asbestos-related diseases.
- Visit by Professor Margaret Becklake, International Asbestos Authority, to advise on registers.
- 1995** *Asbestos Exposure and Disease Notes for Medical Practitioners* published by Department of Labour.
- 1995/96** The third Annual Report confirms 535 cases.
- 1996** *The Management of Asbestos in the Non-Occupational Environment* published by Ministry of Health.
- 1996/97** The fourth Annual Report of the National Asbestos Register notes an increase in occurrence of mesothelioma cases.
- 1997** The Asbestos Medical Panel published its first report based on a review of asbestos cement-exposed workers on the Exposure Register. 2257 notifications were reviewed. A positive finding was a non-specific association between cumulative asbestos exposure and a questionnaire diagnosis of asthma.
- 1998** Health and Safety in Employment (Asbestos) Regulations 1998 promulgated.
- 1999** Visit by Professor Corbett McDonald and Professor Alison McDonald to advise on updated information on mesothelioma.
- Asbestos and the New Millennium* published. Prepared by Lois Syret and Ed Grootegoed on behalf of The Asbestos Association of New Zealand.
- 2000** *Mesothelioma in New Zealand*, the second study by the National Asbestos Medical Panel.
- Increased Mesothelioma Incidence in New Zealand*. T Kjellstrom, P. Smartt. NZMJ, November.

*Lung function changes in asbestos-exposed workers with pleural plaques, the third study by the National Asbestos Medical Panel.*

**2001**

*Asbestos New Zealand 2001, Lois J Syret and A.H. Grootegoed.*

Occupational Safety and Health (Department of Labour) Review of *Guidelines for the Management and Removal of Asbestos.*

*Asbestos Exposure in New Zealand. Bill Glass, New Ethicals, August.*

**2003**

*The New Zealand National Asbestos Registers 1992 to 2001, Lessons Learned. Presented at Asbestos Symposium Seoul National University, Korea by Bill Glass.*

The Epidemic of Asbestos-related Diseases in New Zealand. *Int. J. Occup – Environ Health* 10.2 June 2004 Tord Kjellstrom.

**2004**

Mortality and Morbidity in Asbestos Exposure in New Zealand: The Hidden Legacy of Asbestos Exposure. *NZ Med J* 117; 117: 1 P Smartt.

## **PART 2: REVIEW OF ASBESTOS-RELATED DISEASE NOTIFICATIONS**

### **2.1 Summary**

This report reviews 1198 cases that were notified to the National Asbestos Medical Panel between March 1992 and July 2010. They include:

- 224 cases of mesothelioma
- 116 cases of lung cancer
- 269 cases of asbestosis
- 589 cases of pleural abnormalities.

The number of lung cancer cases reported is roughly half of the number of mesothelioma cases. This suggests that the taking of a lung cancer history is still dominated by the smoking factor, and that occupational factors are downplayed.


The transfer of asbestos from workers to the home is another emerging feature of asbestos-related disease in New Zealand. Family members are presenting with pleural changes or, rarely and tragically, mesothelioma.

Mesothelioma notifications remain high, reflecting exposure in the 1960s and 1970s, and this trend is likely to continue for some years to come.

## 2.2 Results

Figure 2 shows the distribution of the four main diagnostic categories:


- lung cancer
- pleural disease
- asbestosis
- mesothelioma.



**Figure 2: Categories of Disease 1992-2010**

What is noticeable is that pleural disease is the main category, with lung cancer clearly under-represented when compared with mesothelioma.

Figure 3 looks at occupations for the total number of notified asbestos disease cases during the period.



**Figure 3: Notified Asbestos Disease by Occupation 1992-2010**

It is clear that carpenters, plumbers and electricians are together responsible for 67 percent of all cases. These 'all purpose' construction workers are an occupational category at risk particularly because, unlike asbestos cement workers, they are not always seen as being at obvious risk.

The non-occupational category refers to cases where an individual's exposure was not work-related. This includes all cases resulting from secondary or environmental exposure, such as children brought up in the home of an asbestos worker, and people who washed asbestos-contaminated clothes. The 'no known exposure' category refers predominantly to mesothelioma cases where conclusive exposure histories were not available. (Note: the time delay is often 40-50 years from exposure and has either been forgotten, never recognised, or not known by the surviving family member).

### 2.3 Mesothelioma


Mesothelioma - a rare cancer of the pleural membranes on the surface of the lungs - is strongly related to asbestos exposure. The panel reviewed 224 cases of mesothelioma, of which:

- 215 were Caucasian
- 6 were Maori
- 3 were identified as "Other"
- 210 were males, 14 were females
- the mean age at diagnosis was 67 years (range 35-85)
- the mean number of years since first exposure was 45 (range 12-74)
- the mean exposure index was 178 (range 1-780)

- there were 19 current smokers, 125 ex-smokers and 59 never-smokers (information for 21 cases was unavailable).


Asbestos processors, plumbers/fitters/laggers, and carpenters/builders, accounted for over 60 percent of all registered cases.

It has been noted that an asbestos exposure history may be lacking with mesothelioma cases. Our experience suggests that with patience and recognition of the range of likely exposures, it is often possible to obtain evidence of asbestos exposure. In one case the disease developed in a middle-aged woman living in a small rural town. It was revealed that as a teenager she had washed the clothes of an older brother who had been a railway workshop apprentice. Asbestos lagging was used in the repair and maintenance of the boilers, and apprentices were known to use asbestos in 'snowball fights'.



**Figure 4: Distribution of Mesothelioma by Occupation**

## 2.4 National Cancer Figures for Mesothelioma



**Figure 5: Number of Cases of Mesothelioma in New Zealand from 1954 to 2007**

Over the period of 1954 to 2007, a total of 983 cases of mesothelioma have been registered. Figure 5 shows that the total number of cases continues to rise and in 2005 it exceeded 100 for the first time. However, in 2006 the total was 80, and 91 in 2007.

Mesothelioma is very much a disease of old age as Table 7 illustrates, with over 50% of cases occurring to people aged 70 or over.

Age Group	Gender		Total
	Male	Female	
Under 20	0	0	0
20 to 44	7	11	18
45 to 69	404	56	460
70+	436	69	505
<b>Total</b>	<b>847</b>	<b>136</b>	<b>983</b>

**Table 7: Mesothelioma Occurrence by Age Range**

Of the 983 cases, 136 occurred to women and 837 to men. As women are seldom employed directly in the asbestos-exposed workplaces, their exposure could be as a result of "secondary" exposure to dust brought home from work on the hair and clothes of family members.

## 2.5 Lung Cancer


Lung cancer is a cancer of some of the cells in parts of the lung, usually beginning in the lining of the airway.

The panel reviewed a total of 116 cases of lung cancer, of which:

- 110 were Caucasian
- 4 were Maori
- 1 was from a Pacific Island
- 1 was identified as "Other"
- 114 were males, 2 were females
- the mean age at diagnosis was 69 (range 42-86)
- the mean number of years since first exposure was 46 (range 17-63)
- the mean exposure index was 165 (range 12-565)
- there were 25 current smokers, 81 ex-smokers, 7 never smokers and 3 unknown.

Lung cancers are classified according to the type of cell affected. Histological classification revealed 52 cases of squamous cell carcinoma, 29 adeno, 15 small cell, 4 undifferentiated, 1 bronchiolar-alveolar, 4 large cell and 11 cases where classification was not stated. In addition, the tumour sites were as follows: 46 upper lobe, 42 lower lobe, 11 middle lobe, and 17 not stated.

The occupational distribution of lung cancer follows a similar pattern to both pleural plaques and asbestosis, in that plumbers, fitters, carpenters and asbestos processors account for most of the cases. See Figure 6: Distribution of Lung Cancer by Occupation.




**Figure 6: Distribution of Lung Cancer by Occupation**

## 2.6 Asbestosis


Asbestosis is a fibrotic or scarring disease of the lung tissue. The disease develops slowly over many years from initial exposure. It can continue to develop after exposure to asbestos has ceased.

The panel reviewed a total of 269 cases of asbestosis, of which:

- 265 were Caucasian
- 2 were Maori
- 2 were from a Pacific Island
- 253 were males, 16 were females
- the mean age at diagnosis was 68 (range 37-86)
- the mean number of years since first exposure was 43 (range 15-71)
- the mean exposure index was 180 (range 10-720)
- there were 21 current smokers, 192 ex-smokers and 49 never smokers (accurate smoking histories were not available in 7 cases)
- radiological changes showed 154 cases with pleural plaques and/or pleural thickening.



**Figure 7: Distribution of Asbestosis Notifications by Occupation**



**Figure 8: ILO Grading of Asbestosis Cases (n = 150)**

Of the 269 asbestosis cases, 150 were categorised by ILO classification, and others were categorised on the basis of CT, HRCT or pathology, where available. With the recent trend to use HRCT, categorisation by ILO classification is less frequently done.

### **2.6.1 Definition of Asbestosis used in the Register**

An important issue with this disease is ‘what criteria constitute a diagnosis of asbestosis?’ The main point of discussion is the difference between a clinical diagnosis of asbestosis, and a diagnosis suitable for use in a national database where the inclusion of patients with early disease is desirable.

The definition of *Gilson*<sup>a</sup> in his review of asbestosis-related lung conditions in the ILO encyclopaedia has been chosen by the panel and is as follows:

- (a) a history of significant exposure to asbestos dust rarely starting less than 10 years before examination;
- (b) radiological features consistent with basal fibrosis (1/0 and above, ILO 1980);
- (c) characteristic bilateral crepitations; and
- (d) lung function changes consistent with at least some features of the restrictive syndrome.

*Gilson* notes not all criteria need to be met in all cases, but that (a) is essential, and (b) should be given greater weight than (c) or (d). However, occasionally (c) may be the sole sign. Further, he notes that although the restrictive syndrome is the most common pattern (about 40%), in about 10% of cases airway obstruction is the main feature, while in the remainder a mixed pattern is seen.

<sup>a</sup> Gilson JC: *Asbestosis, Encyclopaedia of Occupational Health and Safety, 1983 3<sup>rd</sup> edition, vol 1 pp 187-191*

Of the 269 asbestosis cases:

- all had a significant exposure history, with a mean exposure index of 180 (range 10-720)
- mean latency was 42 years, with a range of 15-71 years
- most cases had an ILO rating 1/1 or greater where this rating was used, although this criterion is now seldom used
- detailed clinical examination results were not always available from the records, thus the presence of crackles was not measurable
- lung function changes are recorded in the register based on the availability of data either from respiratory laboratories, respiratory physicians, or occupational health nurses
- additional information from HRCT scanning has led to the recognition of some cases of asbestosis not covered by the *Gilson* criteria. As noted, HRCT diagnosis is now the norm.

The lung function test numbers in the report confirm the classical restrictive lung function pattern picture does not dominate, with obstructive, mixed, and normal patterns occurring.

## 2.7 Pleural Abnormalities

Pleural abnormalities include pleural plaques, diffuse pleural thickening, chronic fibrosing pleuritis and pleural effusions. It does not include pleural disease occurring together with mesothelioma, lung cancer or asbestosis.




Figure 9: Distribution of Pleural Abnormality Notifications by Occupation

Of the 589 cases reviewed:

- 577 were Caucasian
- 8 were Maori
- 4 were from Pacific Islands
- all but 10 were males
- the mean exposure index was 162 (range 6-708)
- there were 59 smokers, 349 ex-smokers and 148 never smokers (accurate smoking histories were not available in 35 cases).

## **2.8 Discussion**

Information recorded in the Disease Register under-estimates the total burden of asbestos-related disease in New Zealand. This is a consequence of the voluntary nature of the Register, lack of understanding of work as a factor in disease causation by the medical profession, and failure by the Cancer Registry to code occupation in their database. However, the Register continues to serve a useful purpose. There is now greater awareness of the work factor in disease than in 1992 when the Register was established. The Department of Labour now has greater commitment to the importance of occupational illnesses; the Accident Compensation Corporation employs a greater number of occupational doctors, and there are increasing numbers of occupational nurses and safety officers in the private sector.

The Registers, part of the wider Notifiable Occupational Disease System (NODS) operated by the Department of Labour have, in the view of the medical panel, played an important part in encouraging these developments.

### **2.8.1 Pleural Plaques**

One of the aims of the medical panel was to confirm the view that pleural plaques were not just a marker of exposure, but represented a disease state. The Department of Labour publication *Lung Function Changes in Asbestos Exposed Workers with Pleural Abnormalities* in 2000 indicated a clear dose response pattern, including a reduction of FVC and FEV1 with increasing asbestos exposure, independent of smoking habit.

Reference was made earlier to the impact of an asbestos-related occupation on the health of a worker's partner and children. Two cases of pleural plaques notified in the last few years illustrate two women developing widespread plaques in their early 70s. Their only exposure was to asbestos dust brought home on their husbands' clothes which they washed. In one case the husband was an asbestos sprayer; the other, a carpenter.

### **2.8.2 Asbestosis**

The increasing use of HRCT has resulted in the identification of minor degrees of asbestosis often with few, if any, symptoms and no disability. It is possible that these individuals will have a better long-term outlook, although this is not yet established.

### **2.8.3 Lung cancer**

The contribution of occupational asbestos exposure to the causation of lung cancer is well recognised as being under-estimated, and over-attributed to smoking among workers exposed to asbestos. One approach to this issue is to determine the ratio between mesothelioma and lung cancer on the grounds that most mesotheliomas are diagnosed and the majority are regarded as being caused by asbestos exposure at work. Various estimates of such a ratio have been suggested and, as noted by *Kjellstrom*<sup>b</sup> can range from 1 to 10. Even if the lower ratio of 1:2 is taken – based on the mesothelioma cases diagnosed over 1994-2005, for example – some 1,594 cases of lung cancer due to asbestos exposure would have occurred, or approximately 145 a year. It is likely that this figure could be even higher.

### **2.8.4 Mesothelioma**

Reported cases of mesothelioma have continued to rise in New Zealand over the past decade as was shown in Figure 5, and based on the New Zealand Cancer Registry. It is of interest to note the mean exposure index for mesothelioma of 152 - as recorded by the panel - is not dissimilar to exposure indices for pleural plaques (162), lung cancer (162) and asbestosis (180). In other words, mesothelioma, like other asbestos-related conditions, is in general dose dependent.

### **2.8.5 Chronic Obstruction Pulmonary Diseases (COPD) and asbestos exposure**

These conditions are now being recorded if present in individuals with an asbestos-related disease, as well as in those asbestos-exposed workers who have no confirmed asbestos-related lung or pleural disease. Over the past years 33% of the 85 cases of asbestos-related disease also had COPD, 40% among cases of pleural plaques, 45% among asbestosis cases, 80% among lung cancer cases and 0% among cases of mesothelioma. In addition, ten cases that were referred to the Panel because of asbestos exposure but without classical asbestos-related conditions, had COPD.

---

<sup>b</sup> Kjellstrom T: *The Epidemic of Asbestos-Related Diseases in New Zealand*. International Journal Occupational - Environmental Health 2004, 102 June

## **PART 3: NON-ASBESTOS OCCUPATIONAL RESPIRATORY DISEASE**

There has been little emphasis on the importance of occupational asthma, silicosis and work-related chronic obstructive respiratory disease in New Zealand until recent years. However, there is a historical base to non-asbestos occupational respiratory disease as referenced below:

**1938**      *Silicosis: Report of Interdepartmental Committee*, Bulletin No 57, Department of Scientific and Industrial Research, Wellington<sup>1</sup>.

This report noted that since the 1915 Miners' Phthisis Act, 1,576 miners had been granted pensions as a result of silicosis; of those, 1,508 were described as gold miners and 68 as coal miners.

**1940s**      A record of tuberculosis morbidity in hospital nurses recorded in the book *Challenge for Health* by Dr F.S. McLean<sup>2</sup> showed a morbidity rate per 1,000 nursing staff which ranged from 8.9 to 20.3, averaging 16.5 over the years 1943-1949.

**1961**      The *Grey Valley Survey*<sup>3</sup> in which Dr Francis de Hamel surveyed 1,524 miners and ex-miners in the coal industry during 1958 revealed only 32 cases of pneumoconiosis, not all of whom had worked in New Zealand coal mines.

**The 1990s**      This period saw a significant output of publications on non-asbestos occupational respiratory disease, to a large extent due to Professor Neil Pearce of the Centre for Public Health Research at Massey University, together with Dr David Fishwick and Lisa Bradshaw. Over this period occupational asthma was investigated nationally<sup>4</sup> among farmers<sup>5</sup>, sawmill workers<sup>6</sup> and aluminium workers<sup>7</sup>; at the same time respiratory symptoms, lung function and dust levels were measured in the wood industry<sup>8,9</sup>, plywood mill workers<sup>10,11</sup>, hairdressers<sup>12</sup>, mussel openers<sup>13</sup>, welders<sup>14,15,16,17,18,19,20,21</sup> and quarry workers<sup>22</sup>.

**2000**      In 2000 Dr Jeroen Douwes joined the Centre, accelerating the interest in sawmill workers, wood dust and respiratory effects.

### **3.1 Occupational Asthma**

In the Colt Lecture given by Anthony Newman Taylor at the *Ninth International Symposium on Inhaled Particles* at Cambridge University in 2001<sup>23</sup>, Professor Taylor noted some key points about this disease:

- asthma is the most prevalent cause in the United Kingdom, and probably in the western world, of respiratory ill health during working life;
- asthma consists of variable airflow limitation, reversibility and hyper-responsiveness of the airways;
- asthma can be induced by direct toxic damage to the lining of the airways (irritant inducers) or the RADS phenomenon, or as a result of a specific

hypersensitivity response to inhaled proteins and low molecular weight chemicals;

- asthma can be incited in persons who already have hyper-responsive airways when exposed to pollutants in the workplace air;
- evidence is strong that the risk of developing occupational asthma is determined more by the level of exposure to the risk factor than by individual susceptibility;
- asthma induced by work exposures carries a high risk of progression to chronic asthma; and
- there is evidence that the extent of occupational asthma can be reduced by reducing the airborne concentration of the causative agents.

The current programme of the Department of Labour to investigate dirty workplace air is a practical response to the accumulating evidence relating to the causes of work-related asthma.

In New Zealand a debate frequently occurs regarding work-aggravated asthma (WAA), and work-induced asthma (occupational asthma), largely related to whether a compensation claim is accepted or not. While this debate may be relevant to the issue of compensation, it is clearly not relevant to reducing and minimising the impact of asthma at work. This is because both work-aggravated and work-induced asthma are a consequence of exposure to dirty workplace air, and the outcome of both types of asthma is seen in lost time from work. The possibility is that what begins as asthma may in time, if exposure continues, lead to a lack of reversibility and the development of chronic obstructive respiratory disease.

In confirming a case of occupational asthma the Panel requires the following:

- a diagnosis of asthma;
- asthma occurring for the first time at work (work-induced asthma) or asthma made worse at work (work-aggravated asthma);
- a work history of exposure to an asthmagen or to a substance that can irritate the respiratory tract;
- supportive peak flow records indicating a temporal relationship between work and symptoms/peak flow; and
- evidence of reversibility.

### **3.2 Occupational Asthma Case Studies**

#### ***Case 1: A MIG welder with chronic bronchitis***

This man, a non-smoker with no history of childhood asthma or hayfever, began employment as a farm worker. He subsequently worked in the freezing industry and in both occupations suffered no respiratory symptoms.

Over the past eight years he has worked as a welder. Early in this occupation he had a severe episode of metal fume fever when welding galvanised steel. Within a year or two he developed symptoms of cough and black sputum which worsened with time, and also reported that he would get short of breath when moving quickly at work – but not away from work. Fellow workers noticed that

he had a wheeze. His symptoms improved when he began to wear full respiratory protection. He mostly worked with MIG welding.

He was investigated by a respiratory physician with CT, lung function testing, hypertonic saline challenge test, and an exhaled nitric oxide test.

The results showed minor scarring at both lung apices, mild bronchial wall thickening most marked in the mid and lower zones, and more so on the right. There was mild bronchial dilatation most marked in the middle lobe and right lower lobe. With the right lower lobe a number of airways were borderline to mildly bronchiectatic. There were several indeterminate pulmonary nodules in the right lung.

Spirometry showed an FEV<sub>1</sub> 3.98 (94%) predicted, FVC 6.01 (111% predicted) with a ratio of 66%. There was no significant improvement post bronchodilator.

A hypertonic saline challenge report rated spirometry as acceptable and repeatable. Baseline spirometry results indicated a mild obstructive pattern. There was no evidence of significant bronchial hyper-responsiveness.

An exhaled nitric oxide test was reported as normal.

The respiratory physician concluded, given there is no evidence of chronic bronchitis with borderline bronchiectasis on the CT with no other evidence of pulmonary fibrosis, the most likely diagnosis is chronic bronchitis secondary to MIG welding.

### ***Case 2: A spray painter with acute irritant bronchitis***

This man, a non-smoker, used an isocyanate-based paint to spray inside a closed 20-foot shipping container. This container served as a painting booth. The report noted that the extraction fan was not extracting properly and the air line diameter size was incorrect.

It is also reported that prior to this episode he had noted a cough when mixing the paint.

His symptoms were cough, phlegm, wheeze, chest tightness, and shortness of breath.

He was seen by a respiratory physician who noted he did not have asthma as a child. It is recorded that he developed a facial rash at the time. His spirometry two weeks after exposure showed a peak flow of 289 litres/minute, a FEV<sub>1</sub> of 3.6 litres, and FVC of 4.2 litres. Subsequently his peak flow rose to 550 litres/minute and his spirometry FEV<sub>1</sub> 3.92 litres and FVC 4.37 litres – a normal ratio. A subsequent methacholine challenge was reported as negative some six months after the reported episode.

### 3.3 Metal Fume Fever

This condition which commonly occurs to welders cutting or welding galvanised steel, as well as less commonly to workers exposed to other freshly formed metal oxides produced during high temperature processes.

Metal fume fever is usually a self limiting systemic condition occurring four to 12 hours after exposure and characterised by fever, muscle aches and pains, malaise, cough and a metallic taste in the mouth. Recovery usually takes 24 hours and unless the doctor asks about a patient's occupation it is usually diagnosed as the common "flu".

In 2004 the *Journal of Occupational and Environmental Medicine* published a paper on the inflammatory responses and oxidative stress in a group of automotive welders in Taiwan. The main findings of this relatively small cross-sectional study were:

1. higher values for average urine zinc, nickel and copper in full time welders than in a control group;
2. a correlation between average urine zinc, nickel and copper levels and hours of welding per week;
3. a finding that 17.7% of welders had experienced flu-like symptoms in the course of their career;<sup>24</sup>
4. urine zinc levels were associated with white blood count, neutrophil count, eosinophal cancer and inter leukin-6, suggesting the inflammatory response might be mediated by cytokines;
5. the results confirm other studies that welding fume exposure is associated with systemic inflammatory responses as indicated by increased levels of white blood cells and neutrophils while the raised eosinophal levels might suggest an allergic mechanism where local injury results in inflammation with the release of a histamine-like substance; and
6. it was also noted that smoking was significantly associated with urine zinc and nickel levels.

### 3.4 Chronic Obstructive Pulmonary Disease

Chronic obstructive respiratory disease (CORD), or Chronic Obstructive Pulmonary Disease (COPD), as it is now more commonly referred to, "is the fourth leading cause of death worldwide"<sup>25</sup>. It is defined as a condition with airflow limitation which is not fully reversible, is progressive and is associated with an abnormal inflammatory response of the lungs to noxious particles or gases. Historically, and still, the major cause is cigarette smoking. However, there is increasing evidence indicating that exposure to dusts, gases, and fumes at work are linked to the development of COPD. As a result, it is now recognised as an occupational disease in certain situations, with likely additive effects occurring between smoking and some workplace exposures.

Contaminants of air associated in studies with work-related COPD, include welding fumes, silica, coal, oil mist, Portland cement, cotton, grain and wood dusts<sup>26</sup>.

Diagnosis of COPD requires both a clinical and occupational approach and as far as the latter is concerned, a careful occupational history. That is, a chronological list of jobs, what the job entailed, which respiratory exposures occurred, to what extent and for how long.

As a consequence, the panel is now not only recording the presence of COPD in asbestos diagnosed cases, but also in asbestos exposed and non-asbestos exposed cases.

### **3.5 Silica and Silicosis – One Agent, Many Outcomes**

Silica is the most common element found in the earth's crust. When combined with other substances, silica is relatively harmless. Silicosis is a serious lung condition described by the ILO as an accumulation of crystalline silica dust in the lungs and the tissue reaction to its presence. It is caused by inhaling airborne crystalline silica dust in high concentrations over a period of time.

In a review of the outcome of silica exposure, Steenland<sup>27</sup> noted "evidence in recent years indicates that silica causes lung cancer, and probably renal disease, in addition to its well-known relationship to silicosis" and, as indicated in the previous section, it can also result in Chronic Obstructive Pulmonary Disease.

While silicosis is rarely diagnosed in New Zealand, there is evidence that most New Zealand rocks contain some quartz, and with erosion and quarrying the quartose sands tend to contain a higher content of quartz than the parent material<sup>28</sup>. This was re-affirmed in an extractive industry study where the dust measurements carried out by the Department of Labour showed that in 13% of the air samples collected, levels of respirable quartz exceeded 0.2mg/m<sup>3</sup> (the current New Zealand [Workplace Exposure Standard](#)).

### **3.6 Indoor Air Quality (IAQ)**

An article on particle emission characteristics of office printers<sup>29</sup> revealed that among 62 printers investigated, while 60% did not emit submicrometer particles, 40% did, among which 27% were high emitters. However, there is no data to link any harm to these exposures. Office-related respiratory conditions have been recognised for some years with pollutants entering the office environment from outdoor air, as well as pollutants arising from within the office environment. These include organic vapours (Volatile Organic Compounds, or VOCs), fibres and inorganic gases.

With increasing numbers of office workers, attention to indoor air quality is as important as attention to factory and outdoor (quarries, construction) air. The article makes the point that by identifying the high emitters and recognising that emitting rates are printer-type specific and are affected by toner cartridge age, such contaminants of indoor air "can be reduced by a proper choice of printers".

### **3.7 Work-Related Occupational Respiratory Disease**

In an editorial in *Occupational Environmental Medicine*, *Kreiss*<sup>30</sup> looked into the future. She noted recent, "new", occupational respiratory diseases such as flock workers' lung - an interstitial lung disease - hypersensitivity pneumonitis associated with bio-contaminated synthetic metal working fluids, asthma associated with 3-amino-5-mercapto-1, 2, 4-triazole (AMT) in herbicide manufacture, and bronchiolites obliterans from flavouring chemicals used in popcorn plant workers.

*Kreiss* wrote of barriers to the recognition of new respiratory diseases, including:

- clinicians too readily attributing lung disease to smoking;
- the lack of work-related symptoms in some diseases, i.e. occupational industrial diseases;
- long latency of some diseases; and
- lack of knowledge by physicians of patients' workplaces and hazards.

With regard to the recognition of new diseases, she instanced the value of a national state-based surveillance system for occupational asthma.

As to the future and emerging causes of occupational lung disease, *Kreiss* cites asthma related to damp offices. "Building dampness can be considered an emerging cause of occupational lung disease because we are still unsure of causal agents ..."

In addition, *Kreiss* raises the issue of COPD and states: "Although 80% of COPD is attributable to smoking, most of the remainder are likely to have occupational causes." She goes on to say that "control recommendations can be made, even when such relations are based on surrogate exposures and before specific aetiological agents or safe levels of exposure are known".

### **3.8 Other asbestos issues**

#### ***3.8.1 Floor sanders***

In 1992, an audit of floor sanders and their working practices was carried out in Christchurch. This audit revealed a number of features:

- a failure to prevent spread of dust to other rooms;
- lack of a thorough 'clean up' after sanding;
- poor respiratory protection for sanders;
- sanders taking asbestos dust home on their work clothes;
- transfer of dust from sanding machines to vehicles;
- careless disposal of sanding dust; and
- a general lack of understanding about asbestos.

Overseas studies confirmed the New Zealand experience, with high levels of asbestos dust measured in rooms during the sanding process.

### **3.8.2 Asbestos removal workers**

This group of workers was very active in the late 1980s and the 1990s. Although the Department of Labour published the *Guidelines for the Management and Removal of Asbestos (1995)*, undoubtedly a proportion of such workers would have been exposed to significant amounts of asbestos for short- or long-term periods, and some would have transported asbestos dust from work to home on their clothes, boots or body. The National Exposure Register does not identify this group specifically, but they could be the group at most risk, and be a source of asbestos disease in the decades ahead.

### **3.8.3 Brake lining repair workers**

Some brake linings still contain asbestos. Such work is often carried out in designated 'small workplaces'. As such they are less likely to be inspected, and more likely to have inadequate local exhaust ventilation. The work is intermittent and the dose may be long-term and low-level in contrast to demolition workers (high-level and short-term). Nevertheless these workers comprise an ongoing 'at-risk group'.

### **3.8.4 Building maintenance workers**

It is of significance that Great Britain's Health and Safety Executive (HSE) has published two related guidance booklets. The first, *Introduction to Asbestos Essentials* (2001), is specifically aimed at building maintenance workers.

The second, *Asbestos Essentials Task Manual* (2008), is aimed at any worker who may come into contact with asbestos in the course of their work. Such workers include electricians, plumbers, computer installers, telecommunication engineers and others.

### **3.8.5 Demolition workers**

The process of demolition is often carried out over a weekend, utilising casual labour. The presence of asbestos is not necessarily determined prior to a demolition, and as a consequence no knowledge of exposure occurs.

## **APPENDIX 1**

### **Contact details for the Asbestos and Occupational Respiratory Disease Registrar**

Asbestos and Occupational Respiratory Disease Registrar  
Department of Labour  
PO Box 3705  
Wellington 6140

### **Members of the National Asbestos Medical Panel**

W. Glass ONZM, MBChB, DPH, DIH, FFOM, FAFOM (Hon), FFOM (I) (Convenor).

R. Armstrong MBChB (Hons), FRCP, FRACP.

D. Jones MBBS, MRCP (UK), FRACP.

A. Brant MBChB, FRACP, MBA, PhD.

### **Obituary – A H Grootegoed**

The Asbestos Panel noted with sadness the passing of Mr A H (Ed) Grootegoed in May 2010. He was the Chairman of the Asbestos Diseases Association of New Zealand (Inc). Ed did much to highlight the asbestos tragedy in New Zealand, in particular his concern at the dumping of asbestos and rebuilding on old dump sites.

In 2008, Ed received the Air New Zealand Lifetime Achievement Award for his work with asbestos concerns.

## **APPENDIX 2**

### **The National Asbestos Registers**

The National Asbestos Registers were established in March 1992 in line with recommendations made to the Minister of Labour by the Asbestos Advisory Committee.

#### ***Formation of the Asbestos Advisory Committee***

The Asbestos Advisory Committee was established in October 1990 to report to the Minister of Labour on issues relating to health effects and use of asbestos in New Zealand, adequacy of controls and legislation, and clarification of the legal entitlements available for affected workers. This followed increasing public concern about past and present effects of asbestos on workers, former workers and their families.

#### ***Establishment of the National Asbestos Registers***

Recommendation 4 of the Report of the Asbestos Advisory Committee to the Minister of Labour advised:

*“That an asbestos medical register be established for people who have been significantly exposed to asbestos. OSH should be the organisation responsible for establishing, maintaining and funding the medical register.*

*The medical register should be in two parts:*

*Part 1 – Those notified as having been exposed to asbestos;*


*Part 2 – Those notified as having an asbestos-related disease.*

*The system should allow movement of the name of a registered person from part 1 to part 2 of the register when indicated.*

*Notifications to part 1 of the medical register were to be made by those who felt they had been exposed to asbestos, or by people acting on their behalf (and following consultation), such as an employer, union official, relative or friend.*

*Notification to part 2 of the medical register would be done by medical practitioners.”*

A Notifiable Occupational Disease System (NODS) was established in 1992 and the Asbestos Registers have been incorporated in that scheme. This was in accordance with recommendation 5 of the Asbestos Advisory Committee.



**Figure 10: Asbestos Exposure Register Process**

The Department of Labour, in association with Electricorp Production Ltd, undertook an extensive advertising campaign in March and April 1992. Advertisements were published in all the major newspapers and several trade magazines.

The interest generated as a result of this campaign has ensured a high response rate for the Exposure Register. Notifications have been made by individuals, trade unions, occupational health nurses, doctors, the Asbestos Diseases Association of New Zealand and by some larger companies.


Notifications are directed to branch offices of the Department or directly to the Registrar.

In recommendation 4, the committee envisaged that people wishing to be recorded on the asbestos exposure register would have their exposure assessed at an office of the Department. Only those people who were judged as having had 'significant exposure' would then be recorded on the register. However, the huge response from individuals exposed made it impractical to screen registrants in this fashion, although this has since been done.

Once a person notifies the Department of Labour that they have been exposed to asbestos, an asbestos exposure registration form is sent. The form collects information about the individual, their work exposure to asbestos and the state of their respiratory health.

When the completed form is returned to the registrar, the details are recorded on a database. The individual is then sent a copy of *Asbestos – A Deadly Dust*, which is dedicated to asbestos and its associated health problems. If the person indicates that they have a family doctor, the doctor is informed their patient has

been included on the Asbestos Exposure Register, and is sent a copy of the Department's booklet *Asbestos Exposure and Disease: Notes for Medical Practitioners*.



**Figure 11: Asbestos Disease Register Process**

Tenders for the National Asbestos Medical Panel were called for in 1991. A tender was accepted on 31 October 1991. The successful tender came from the group listed below:

Dr R. Armstrong, Professor R. Beasley, Dr J. Crane, Associate Professor W. Glass, Dr D. Jones, Dr N. Pearce.

Professor Beasley retired upon his appointment as Professor of Medicine at the Wellington Clinical School. Dr Crane joined the National Occupational Asthma Panel. Dr D. Fishwick joined the panel in 1997. He was subsequently appointed to a position in the United Kingdom. In 2008, Andrew Brant joined the Panel.

The first meeting of the panel was held in February 1992. Professor Glass was nominated as the panel's convenor.

The following members were appointed to the National Asbestos Radiological Panel: Dr Paul White, Dr George Foote and Dr Graeme Anderson.

The Registrar from 1991 to 1996 was Craig Eades. From 1996 to 1998 the position was held by Nicola Holden, and later in the year by Andrea Eng until 1999 when Louisa Thomas was appointed.

In 2002 Jenny West was the Registrar, with Dougal McNeill Registrar in 2004, Gail Abel in 2006, Justine Ward and Christy He in 2007 and Anne Thomas in 2008.


### ***Processes for registering people***

Notifications for the register come from two major sources. The first is from doctors whose patients have been diagnosed or are suspected of having an asbestos-related disease. The second source of notification is from individuals. Once a notification is made to the Registrar, and consent gained from the person concerned, relevant medical records and a full occupational history are obtained.

### ***Data collection***

The data collected includes a medical history, occupational history, chest x-ray, CT scan where available, lung function tests and pathology reports. On notification being received by the Registrar:

- An occupational health nurse visits the patient and conducts a health interview, a detailed occupational and social (including smoking) history.
- Relevant medical reports are obtained from general practitioners and physicians.
- A PA chest x-ray is obtained and read by a radiologist according to ILO (1980) guidelines. CTs are used where available, and on occasions requested.
- Lung function data is obtained from physicians' reports or requested from respiratory laboratories. Where this is not possible, results are obtained from a test carried out by an occupational health nurse, using a portable spirometer.
- Pathology and post-mortem reports are reviewed where available.



This is calculated for each job and total exposure is the sum of each D.

**Figure 12: Exposure Index Calculation Formula**

### ***Data assessment***

The National Asbestos Medical Panel reviews the information obtained, calculates an exposure index (see overleaf) and correlates the medical data.

### ***Exposure Index***

An exposure index (D) was calculated from the product of years of asbestos exposure (A); intensity of exposure, using a 1-5 grading according to job category (B); and degree of exposure, using a 3-point grading (C).

Guidelines for calculating this index are shown below.

- A** = Total years of exposure in any one job
- B** = Job intensity as follows:  
Mining, milling and processing = 5  
Boiler/lagging, rail carriages, shipyard, spraying insulation = 4  
Asbestos cement products, construction, demolition, removal = 3  
Electrical, friction products = 2  
Loading, driving, environmental = 1
- C** = Degree of exposure (unprotected)  
Continuous (>50% of work) = 3  
Intermittent (20-50% of work) = 2  
Minimal (<20% or occasional) = 1
- D** = A x B x C for each job  
Exposure index = sum of all Ds

### ***Medical data***

Relevant respiratory symptoms and signs are recorded from the medical histories, and lung function data is classified into restrictive, obstructive, mixed or normal. Pathology reports are used to confirm mesotheliomas and classify lung cancers.

### ***Classification of Diagnostic Categories***

On the basis of the foregoing, the cases were placed into a primary diagnostic category of:

- mesothelioma
- lung cancer
- asbestosis
- pleural abnormalities (plaques, diffuse bilateral pleural thickening and effusions)
- other cancers
- obstructive lung disease without x-ray changes.

## APPENDIX 3

### Department of Labour Publications on Asbestos

1. *Talking Asbestos* (brochure) 1987 (Archived)  
<http://www.osh.govt.nz/order/catalogue/archive/talkingasbestos.pdf>
2. *Asbestos Exposure and Disease: Notes for Medical Practitioners.* (Information Bulletin) 1992.  
<http://www.osh.dol.govt.nz/order/catalogue/237.shtml>
3. *Guidelines for the Management and Removal of Asbestos*, 1995.  
<http://www.osh.govt.nz/order/catalogue/pdf/asbmgmt.pdf>
4. *The Epidemiology of Mesothelioma in Historical Context.* J.C. McDonald and A.D. McDonald, 1998. Special Report. (Archived)  
<http://www.osh.govt.nz/order/catalogue/archive/mesothelioma.pdf>
5. *Respiratory Symptoms and Asbestos Dust Exposure.* Occupational Health Report Series: No. 2, 1997.  
<http://www.osh.govt.nz/order/catalogue/pdfs/ohrs002.pdf>
6. *Asbestos: A Deadly Dust. 50 Years of Asbestos Use in New Zealand.* Reprint from *Safeguard* magazine, December 1991.
7. *National Asbestos Registers, Annual Reports: 1992-1993, 1994-1995, 1995-1996, 1996-1997, 1997-1998, 1992-2004, 1992-2005, 1992-2007, 1992-2008* <http://www.osh.govt.nz/order/catalogue/524.shtml>
8. *Respiratory Health and Silica Dust Levels in the Extractive Industry.* Occupational Health Report Series: No. 9, 2003.  
<http://www.osh.govt.nz/order/catalogue/pdfs/ohrs009.pdf>
9. *Report of the Asbestos Advisory Committee to the Minister of Labour.* Department of Labour, 1991.  
<http://www.osh.govt.nz/order/catalogue/archive/asbestosreport.pdf>
10. *Flooring Contractors.* Occupational Health Report Series: No. 4, 2000.  
<http://www.osh.govt.nz/order/catalogue/pdfs/occ4flooring.pdf>

#### ***Out-of-Print Publications (limited availability)***

1. *Mesothelioma in New Zealand* Glass, Bill, Fishwick, D. Armstrong, R. Jones, D. Bradshaw, L.M. Department of Labour, Wellington, New Zealand, 2000.
2. *Lung Function Changes in Asbestos Exposed Workers with Pleural Abnormalities* Armstrong, R et al, Department of Labour, Wellington, New Zealand, 2000.

3. *Recent Advances in Asbestos-Related Disease; Asbestos and other Fibre-Related Diseases of the Lung and Pleura: Distribution and Determinants in Exposed Populations.* Becklake, M.R. Glass, B. Armstrong, R. Eades, C. Department of Labour, Wellington, New Zealand, 1991.

## APPENDIX 4

### References

---

- <sup>1</sup> *Silicosis Report of Interdepartmental Committee* Bulletin no. 57, Department of Scientific and Industrial Research 1937 Government Printer, Wellington.
- <sup>2</sup> McLean, E.S. *Challenge for Health* 1964. Government Printer, Wellington.
- <sup>3</sup> de Hamel, F.A. *The Grey Valley Survey* 1961. Department of Health, Wellington.
- <sup>4</sup> *Occupational Asthma in New Zealanders: A Population-Based Study*. D. Fishwick, N. Pearce et al. *Occupational and Environmental Medicine* 1997; 54: 301-306.
- <sup>5</sup> *Asthma and Allergy in New Zealand Farmers*. M. Kimbell-Dunn, L.M. Bradshaw et al. *American Journal of Industrial Medicine* 1999; 35: 51-57.
- <sup>6</sup> *Asthma and Other Respiratory Symptoms in New Zealand Pine Processing Sawmill Workers*. J. Douwes, D. MacLean et al. *American Journal of Industrial Medicine* 2001; 39: 608-615.
- <sup>7</sup> *Potroom Asthma: New Zealand Experience and Follow-Up*. T.V. O'Donnell, B. Wellford and E.D. Coleman. *American Journal of Industrial Medicine* 1989; 15: 43-49.
- <sup>8</sup> *Worker Exposures to Airborne Dust, Endotoxin and B(1, 30 – Glucan) in Two New Zealand Sawmills*. J. Douwes, D. MacLean et al. *American Journal of Industrial Medicine* 2000; 38: 426-430.
- <sup>9</sup> *Wood Dust Exposure in Small New Zealand Woodworking Companies*. D. McBride, K. Nelse. *The Journal of Occupational Health and Safety – Australia and New Zealand*, 2004; 20: 465-472.
- <sup>10</sup> *Respiratory Symptoms and Occupational Exposures in New Zealand Plywood Mill Workers*. W. Fransman, D. MacLean et al. *American Occupational Hygiene* 2003; 42: 287-295.
- <sup>11</sup> *Pine Dust, Atopy and Lung Function: A Cross-Sectional Study in New Zealand Sawmill Workers*. J. Douwes, D. MacLean, T. Slater, N. Travier, S. Cheng, N. Pearce. *European Respiratory Journal*. 2006; 28: 791-798.
- <sup>12</sup> *Occupational Respiratory Symptoms in New Zealand Hairdressers*. T. Slater, L. Bradshaw et al. *Occupational Medicine* 2000; 50: 586-590.

- 
- <sup>13</sup> *Work-Related Respiratory Symptoms and Lung Function in New Zealand Mussel Openers.* W.I. Glass, P. Power et al. *American Journal of Industrial Medicine* 1998; 34: 163-168.
- <sup>14</sup> *Chronic Interstitial Lung Disease in a Welder of Galvanised Steel.* W.I. Glass, D.B. Taylor, A.M. Donoghue. *Occupational Medicine* 1994; 44: 158-160.
- <sup>15</sup> *Transient Changes in the Pulmonary Function of Welders: A Cross-Sectional Study of Monday Peak Expiratory Flow.* A.M. Donoghue, W.I. Glass, G.P. Herbison. *Occupational and Environmental Medicine.* 1994; 51: 553-556.
- <sup>16</sup> *Chronic Bronchitis, Work-Related Respiratory Symptoms, and Pulmonary Function in Welders in New Zealand.* L.M. Bradshaw, D. Fishwick et al. *Occupational and Environmental Medicine* 1998;55: 150-154.
- <sup>17</sup> *Two Year Follow-Up of Pulmonary Function Values among Welders in New Zealand.* R. Erkinjuntti-Pekkanen, T. Slater et al. *Occupational and Environmental Medicine* 1999; 56: 328-333.
- <sup>18</sup> *Exposure to Fumes in Typical New Zealand Welding Operations.* E. Dryson, D. Rogers. *New Zealand Medical Journal* 1991; 23: 351-258.
- <sup>19</sup> *Respiratory Symptoms, Across-Shift Lung Function Changes and Lifetime Exposures of Welders in New Zealand.* D. Fishwick, L. Bradshaw et al. *Scandinavian Journal of Work and Environmental Health.* 1997; 23:351-358.
- <sup>20</sup> *Changes in Work Practice after a Respiratory Health Survey among Welders in New Zealand.* T. Slater, R. Erkinjuntti-Pekkanen, D. Fishwick, L. Bradshaw, N. Pearce, S. Cheng, H. Armstrong, D. McLean. *New Zealand Medical Journal* 2000 July 28; 113 (1114): 305-8.
- <sup>21</sup> *Respiratory Symptoms and Lung Function Change in Welders: Are they associated with Workplace Exposures?* D. Fishwick, L. Bradshaw, A. Curran, N. Pearce. *New Zealand Medical Journal* 2004 May 7; 117 (1193): U872.
- <sup>22</sup> *Respiratory Health and Silica Dust Levels in the Extractive Industry in New Zealand.* W.I. Glass, D. McLean et al. *Occupational Health Report Series, No. 9* 2003; 25.
- <sup>23</sup> *Asthma and Work.* Anthony Newman Taylor. *Ann. Occupational Hygiene* 2002; 46: 563-574.
- <sup>24</sup> *Inflammatory Responses and Oxidative Stress from Metal Fume Exposure in Automobile Welders.* Luo, Jiin-Chyuan John, Hsu, Kuang-Hung, Shen, Wu-Shiun. *Journal of Occupational And Environmental Medicine* 2009;51(1): 95-103.
- <sup>25</sup> *Occupation and COPD.* D.C. Christiani. *Occupational and Environmental Medicine* 2005; 62: 215.

---

<sup>26</sup> *Occupational Contribution to the Burden of Chronic Obstructive Pulmonary Disease.* John Balmes. *Journal of Occupational and Environmental Medicine* 2005; 47: 154-160.

<sup>27</sup> *One Agent, Many Diseases: Exposure-Response Data and Comparative Risks of Different Outcomes Following Silica Exposure.* K. Steenland. *American Journal of Industrial Medicine* 2005; 48: 16-23.

<sup>28</sup> *Exposure to Alpha Quartz Dust in New Zealand Industries.* E. Nicol. *Occupational Health* 1982; 4:1, 30-36.

<sup>29</sup> *Particle Emission Characteristics of Office Printers.* Congreve H. E. Congrong, L. Morawska, L. Taplia. *Environmental Science and Technology* 2007; 41:17 6039 – 6045.

<sup>30</sup> *Emerging Opportunities to Prevent Occupational Lung Disease.* Kathleen Kreiss. *Occupational and Environmental Medicine* 2007; 64: 499-500.



FOR MORE INFORMATION ON HEALTH AND SAFETY  
VISIT [WWW.DOL.GOVT.NZ](http://WWW.DOL.GOVT.NZ)



PLA 11590 DEC 10

